

Assessment/Release for Return to Play

Patient: School:

DOB: Sport:

PCP:

-

Date of onset of COVID symptoms:

Date of COVID positive test:

Date of resolution of COVID symptoms:

Systemic symptoms for 5 days or more (fever, myalgia, chills, profound lethargy)?: N Y

Hospitalization due to COVID symptoms?: N Y

H/o cardiac abnormalities followed by cardiology?: N Y

Recent symptoms:

Chest pain at rest or with exertion? (not musculoskeletal or costochondritis): N Y

Shortness of breath with minimal activity? (unrelated to respiratory symptoms): N Y

Excessive fatigue with exertion?: N Y

Abnormal heartbeat or palpitations?: N Y

Syncope or near-syncope?: N Y

Normal cardiovascular exam?: Y N

Cardiology referral indicated?: N Y

Cleared for gradual return to sports?: Y N

Signature:

Printed Name:

Date:

Suggested gradual return to sports progression*

(adapted from Elliott N, et al, infographic, British Journal of Sports Medicine, 2020):

Stage 1: 2 days minimum, ≤ 15 minutes, light activity (walking, jogging, stationary bike), no resistance training.

Stage 2: 1 day minimum, ≤ 30 minutes, add simple movement activities (eg. running drills).

Stage 3: 1 day minimum, ≤ 45 minutes, progress to more complex training, may add light resistance training.

Stage 4: 2 days minimum, 60 minutes, normal training activity.

*Patient should be symptom free before progressing to next stage.